

# **The Oxidative Coupling of Aliphatic Aldehydes**

By JOHN C. LEFFINGWELL

*(Research and Product Development Departments, R. J. Reynolds Tobacco Company, Winston-Salem, North Carolina 27102)*

Reprinted from

## **Chemical Communications 1970**

The Chemical Society, Burlington House, London W1V 0BN

## The Oxidative Coupling of Aliphatic Aldehydes

By JOHN C. LEFFINGWELL


(Research and Product Development Departments, R. J. Reynolds Tobacco Company, Winston-Salem, North Carolina 27102)

**Summary** The C-C and C-O oxidative coupling of aliphatic aldehydes is reported.

absorptions at 3.59 and 5.76  $\mu\text{m}$  (saturated aldehyde); n.m.r. ( $\text{CDCl}_3$ ): gem dimethyls at  $\tau$  8.84 and aldehydic proton

MANGANESE DIOXIDE and nickel peroxide are classical mild oxidants for the conversion of allylic and benzylic alcohols into  $\alpha\beta$ -unsaturated aldehydes and ketones.<sup>1-3</sup> Recently we reported the oxidative dehydrogenation of various cyclohexene aldehydes with active manganese dioxide as a convenient preparative procedure for the production of aromatic aldehydes.<sup>4</sup> We now report that  $\alpha$ -substituted aliphatic aldehydes readily undergo  $\alpha$ -hydrogen abstraction and subsequent C-C and C-O dimerization without further oxidation of the -CHO functionality or appreciable formation of  $\alpha\beta$ -unsaturation.<sup>5</sup> Although the oxidative coupling of phenolic materials<sup>6</sup> and aromatic amines<sup>7</sup> has been widely reported, only a few examples of the C-C and C-O coupling of ketones<sup>8</sup> or  $\beta$ -diketones<sup>9</sup> have appeared, and no previous reports of aldehyde couplings are available. The results of our preliminary work indicate that certain aldehydes are in fact easily dimerized by a number of common oxidizing agents, and the products are stable, easily isolable materials of considerable synthetic interest. Among the reagents which are effective for such oxidative aldehyde couplings are activated manganese dioxide, nickel peroxide, and lead dioxide.

For example, when isobutyraldehyde (I;  $\text{R}^1 = \text{R}^2 = \text{Me}$ ) (18 g) in THF (150 g) was continuously refluxed through a modified Soxhlet cup containing activated manganese dioxide (50 g) for 48 hr. ca. 80% of a dimeric mixture of 2,2,3,3-tetramethylsuccinaldehyde (II;  $\text{R}^1 = \text{R}^2 = \text{Me}$ ) [I.r.



resonance at 0.53 in a ratio of 6:1; mass spectrum: molecular ion at  $m/e$  142] and 2-methyl-2-(2'-methyl-1'-propenoxy)propionaldehyde (III;  $\text{R}^1 = \text{R}^2 = \text{Me}$ ) [I.r.: 3.66 and 5.75 (saturated aldehyde) and 5.93  $\mu\text{m}$  (vinyl ether); n.m.r. ( $\text{CDCl}_3$ ): gem dimethyls at  $\tau$  8.75 (6H), allylic methyls centred at 8.45 (6H), olefinic proton at 4.34 (1H), and aldehydic proton at 0.55 (1H); mass spectrum: molecular ion at  $m/e$  142] was produced. The resultant mixture (see Table) was conveniently separated by steam distillation in that the enol ether-aldehyde was rapidly distilled while the tetramethylsuccinaldehyde formed a hydrate obtained by solvent extraction of the steam pot. Approximate ratios of C-C and C-O coupling products for some representative systems are in the Table. Product ratios are virtually identical using nickel peroxide as

Aldehyde	Reagent	Solvent	Coupling product ratios	
			C-C(%)	C-O(%)
Isobutyraldehyde .. .. .	MnO <sub>2</sub>	THF	43	: 57
2-Methylbutyraldehyde .. .. .	MnO <sub>2</sub>	THF	44	: 56
2-Ethylbutyraldehyde .. .. .	MnO <sub>2</sub>	Dioxan-pyridine	12	: 88
Cyclohexanecarboxaldehyde .. .. .	PbO <sub>2</sub>	Dioxan	45	: 55

oxidant; however, low yields are generally obtained with this oxidant.

The C-O dimerization product (III; R<sup>1</sup> = R<sup>2</sup> = Me) from isobutyraldehyde which formally appears to meet the requirements for a potential thermally degenerate system was examined by n.m.r. over a varying temperature range up to 150° without any indication of valence isomerism.

Finally, our experiments in which C-O coupling products of  $\alpha$ -substituted aldehydes are actually isolated offer an alternative explanation for the formation of  $\alpha$ -hydroxyaldehydes in aqueous acid media by one electron oxidants.<sup>10</sup>

(Received, January 21st, 1970; Com. 102.)

<sup>1</sup> S. Ball, T. W. Goodwin, and R. A. Morton, *Biochem. J.*, 1948, **42**, 516.

<sup>2</sup> R. M. Evans, *Quart. Rev.*, 1959, **13**, 61.

<sup>3</sup> K. Nakagawa, R. Konaka, and T. Nakata, *J. Org. Chem.*, 1962, **27**, 1597; K. Nakagawa, R. Konaka, and J. Sugita, *Ann. Report Shionogi Res. Lab.*, 1969, **19**, 141.

<sup>4</sup> J. C. Leffingwell and H. J. Bluhm, *Chem. Comm.*, 1969, 1151.

<sup>5</sup> C. W. Hargis and H. S. Young, *Ind. and Eng. Chem. (Product Res. and Development)*, 1966, **5**, 72.

<sup>6</sup> A. I. Scott, *Quart. Rev.*, 1965, **19**, 1; H. Musso, *Angew. Chem.*, 1963, **75**, 965; J. R. Lewis, *Chem. and Ind.*, 1962, 159; E. McNelis, *J. Org. Chem.*, 1966, **31**, 1255.

<sup>7</sup> O. Meth-Cohn and H. Suschitzky, *Chem. and Ind.*, 1969, 443 and references therein; S. Terabe and R. Konaka, *J. Amer. Chem. Soc.*, 1969, **91**, 5655.

<sup>8</sup> A. Wolf, G.P. 876,237/1953; E. Wolthuis, B. Bossenbroek, G. DeWall, E. Geels, and A. Leegwater, *J. Org. Chem.*, 1963, **23**, 148; H. C. Volger and W. Brackman, *Rec. Trav. chim.*, 1965, **84**, 1233; M. F. Ansell, W. J. Hickenbottom, and P. G. Holton, *J. Chem. Soc.*, 1955, 349; M. S. Kharasch, H. C. McBay, and W. H. Urry, *J. Amer. Chem. Soc.*, 1948, **70**, 1269; J. C. Leffingwell, F.P. 1,544,604/1968.

<sup>9</sup> H.-J. Teuber, *Angew. Chem. Internat. Edn.*, 1969, **8**, 218; O. H. Mattsson and C. A. Wachtmeister, *Acta Chem. Scand.*, 1968, **22**, 79; *Tetrahedron Letters*, 1967, 1855; W. Reusch and R. Starkey, *J. Org. Chem.*, 1967, **32**, 931; H.-D. Becker, *J. Org. Chem.*, 1965, **30**, 989.

<sup>10</sup> W. A. Waters, "Mechanisms of Oxidation of Organic Compounds," Methuen, London, 1964, pp. 87-88; *Progr. Org. Chem.*, 1961, **5**, 1, and references therein.