

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58

65 | Cooling Ingredients and Their Mechanism of Action

John C. Leffingwell

Leffingwell & Associates, Canton, Georgia, U.S.A.

INTRODUCTION

The use of “purified” cooling agents in pharmaceutical and cosmetic preparations only dates back to the late 1880s with the commercial production of menthol from Japanese peppermint (*Mentha arvensis*) oil in Japan (1). The cultivation of peppermint in Japan before the Christian era appears to predate any other country, and menthol is reputed to have been used medicinally for almost as long (2). In the Western world, it was about 1770 that the Dutch botanist, H. David Gaubius, first isolated menthol from the oil of *Mentha piperita* in Utrecht (2,3). Prior to the commercial availability of menthol, the essential oils of peppermint varieties (primarily *M.* and *M. arvensis*) were the sole source for use as cooling minty ingredients. It is significant that at the end of the 18th century only about 900 to 1400 kg of peppermint oils (both *piperita* and *arvensis*) were consumed worldwide (1). By the late 1890s, production had increased to about 175,000 kg (2). In 2007, total peppermint oil production was estimated at more than 26,000,000 kg, with about 21,500,000 kg being the oil of *M. arvensis* (commonly referred to as cornmint oil), which is used mostly for the production of natural *leavo*-menthol (4).

This chapter reviews the use of menthol and new classes of cooling agents that have been discovered since the 1970s. In addition, we briefly touch upon the efficacy of cooling agents as insect repellents. Finally, recent findings on the physiological mechanisms of cold receptors are presented.

COOLING INGREDIENTS

Menthol Background

Before World War II, production of *leavo*-menthol [hereafter referred to as (–)-menthol] was controlled exclusively by Japan and China. In 1939, Japan exported 268,920 kg of menthol, while China’s exports in 1940 were 190,909 kg (1). With the advent of war, shipments to the allied countries ceased and major shortages ensued. While synthetic (–)-menthol could be produced from high citronellal feed stocks (e.g., citronella oil and citronella-type eucalyptus oils), this also was no longer an option. However, Japanese and Chinese immigrants in Brazil rapidly began planting *M. arvensis* for menthol production. In 1941, Brazil produced 5000 kg of menthol, rising to 1,200,000 kg by 1945 (1). By the 1960s, Brazil’s production peaked at about 3,000,000 kg, while about the same time China began supplying menthol again.

During the 1960s, an oversupply of menthol caused the price to fall to as low as \$7.70 to \$8.80 per kilogram, and processors reduced production levels. This ultimately led to worldwide shortages and a price spike as high as \$50 plus per kilogram in 1974 (with similar price spikes now occurring about every 10 years) (3). As menthol is a commodity, it is sometimes subject to financial speculation, which exacerbates price swings.

In 1958, India began expanding plantings of *M. arvensis*, but, until the late 1980s, the quality was highly variable and often had low menthol content. In the 1980s, new strains were introduced that gave improved oil yields and had menthol contents of 75% to 85%. By 1996, India was producing 6000 metric ton of *M. arvensis* oil and had long surpassed China as the major producer of menthol (3). In 2007, it was estimated that India would produce in excess of 20,000 metric ton of this mint oil. While the bulk of current production is used for local menthol crystallization, significant amounts of oil and crude menthol fractions are exported to Brazil, Taiwan, and Japan for further purification. The residual oil left after crystallizing

59 menthol still contains 35% to 45% menthol as well as menthones and other typical mint
 60 components. Much of this oil (commonly referred to as dementholized cornmint oil) is
 61 rectified by distillation and sold for use where normal peppermint oil (ex *M. piperita*) is used
 62 (toothpaste, mouthwash, etc.). In addition, some of this dementholized oil is fractionated to
 63 isolate the menthones (which can be converted by reduction into (-)-menthol) and other
 64 "natural" flavor chemicals.

65 During the 1970s and 1980s, a number of new routes to synthetic (-)-menthol were
 66 developed, only two of which led to long-term commercial success. These processes have been
 67 reviewed by both Leffingwell (5) and Hopp and Lawrence (6). Today, the procedure
 68 developed by Haarmann and Reimer (now Symrise) on the basis of hydrogenation of thymol
 69 to racemic *dl*-menthol followed by selective crystallization of (-)-menthol (via the benzoate
 70 ester) is the major process (7).

Q1

71 The Takasago process uses myrcene as the raw material, which is converted to *N,N*-
 72 diethylgeranylamine and then asymmetrically isomerized via the chiral rhodium (S)-BINAP
 73 (or SEGPPOS) complex to the optically active enamine of citronellal. Hydrolysis yields (+)-
 74 citronellal, which is cyclized to (-)-isopulegol by classical methods. On hydrogenation, the
 75 isopulegol gives (-)-menthol in high optical purity (8,9). An alternative starting material
 76 (instead of myrcene) is isoprene, which can be dimerized to *N,N*-diethylnerylamine. This
 77 material can be converted to (-)-menthol in a manner analogous to the myrcene route using
 78 rhodium (R)-BINAP as the chiral catalyst (10). Reflecting on the Takasago process, Ryoji
 79 Noyori stated in his 2001 Nobel lecture, "This resulted from a fruitful academic/industrial
 80 collaboration..." (11).

81 Clark estimates that 2007 worldwide consumption of menthol from all sources (i.e.,
 82 peppermint oils, natural menthol, and synthetic menthol) is 32,000 metric ton, of which 19,170
 83 metric ton is purified menthol (3).

Q2

84 Table 1 provides our estimate of production in producing countries (or in the case of
 85 Symrise and Takasago, company production of synthetic (-)-menthol).

86 Table 2 provides the 2007 estimated worldwide usage of menthol by consumer product
 87 category—on the basis of Clark's data by region (3).

Q3

90 **Table 1** Worldwide Sources of Menthol (2007)

Source	Metric ton
India (natural)	9,700
China (natural)	2,120
Symrise (synthetic)	3,600
Takasago (synthetic)	1,500
Other synthetic ^a	1,200
Brazil (natural) ^b	450
Taiwan (natural) ^b	300
Japan (natural) ^b	300
Total ^c	19,170

102 ^aOther synthetic includes menthol produced from menthones as well as racemic menthol.

103 ^bPrimarily from *Mentha arvensis* oil or crude menthol ex India (or China).

104 ^cTotal menthol volume based on Clark's estimate (4).

Q11

107 **Table 2** 2007 Estimated Worldwide Consumption of Menthol % by Product Category

Product category	Menthol %
Oral hygiene	28.00
Pharmaceuticals	26.60
Tobacco	25.30
Confectionaries	11.00
Shaving products	7.00
Miscellaneous	2.10

116 *Source:* From Ref. 4.

Q9

117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174

Menthol Chemistry

Menthol is a C₁₀H₂₀O terpenoid alcohol (MW 156.27) with three chiral centers leading to eight possible stereoisomers (4 enantiomeric pairs). The characterization of the stereoisomeric menthols was painstakingly resolved prior to the availability of modern methods by Read (12,13). The structures of the eight enantiomers, with their optical rotations (in ethanol), are shown in Figure 1.

Only the (-)-menthol enantiomer possesses the clean desirable minty odor and intense cooling properties. For example, the (+)-menthol enantiomer is less cooling and possesses a musty off-note odor that is undesirable in most applications. This musty note is also present in racemic menthol (15). The organoleptics and cooling strengths of all of the enantiomers have been reviewed (5,6). Figure 2 provides the cooling thresholds in ppm.




Figure 1 Stereoisomers of menthol. Source: From Ref. 14. Q10




Figure 2 Cooling thresholds (in ppm) (by taste dilution). Source: From Ref. 14.

175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232

Table 3 Major Impurities in Synthetic and Natural Menthols

Major impurities	Synthetic %	Brazil 1 %	Brazil 2 %	China 1 %	China 2 %	India %
Menthone	0.0069	0.0258	0.0258	0.0135	0.0350	0.0295
Isomenthone	0.0000	0.0069	0.0172	0.0052	0.0123	0.0155
Menthyl acetate	0.0000	0.0100	0.0148	0.0014	0.0128	0.0048
Isopulegol	0.0022	0.1868	0.1651	0.1374	0.1914	0.1789
Neomenthol	0.0032	0.0689	0.1339	0.0951	0.0882	0.1079
Neoisomenthol	0.0000	0.0075	0.0459	0.0352	0.0177	0.0368
Isomenthol	0.0299	0.0099	0.0442	0.0296	0.0248	0.0322
Piperitone	0.0000	0.0053	0.0046	0.0018	0.0031	0.0024
Totals	0.0422	0.3211	0.4515	0.3192	0.3853	0.4080

Source: From Ref. 16.

Natural menthol ex *M. arvensis* oil is normally about 99.0% to 99.6% pure, with the remaining impurities being other constituents found in the cornmint oil. While, in most cases, the mint oil impurities contribute a pleasant peppermint aroma, certain impurities, such as mint sulfide, can also impart less desirable and harsh notes. Thus, odor discrepancies often arise when comparing samples from different companies or countries. To overcome such differences, the skilled technician can add a small percentage (e.g., 0.2–0.4%) of terpeneless peppermint oil ex *M. piperita* (or redistilled dementholized cornmint oil), which adds the desirable sweet peppermint top note. Table 3 compares the major impurities present in synthetic menthol and natural menthol samples from major producing areas (16).

Although not generally commercially available, menthol produced from *M. piperita* oil has a sweeter peppermint top note than that produced from cornmint oil (JC Leffingwell, unpublished observations).

Synthetic (–)-menthol is normally about plus 99.8% pure and has less of the minty top note present in natural menthol. Again, this can be adjusted to increase the mint character, if desired, by the addition of a small amount of terpeneless peppermint oils.

Menthol-Related Cooling Agents

Interest in menthol-related cooling agents began in the late 1950s to 1960s when several tobacco companies began to develop various esters as potential menthol release agents (17–19), some of which now appear on the flavor extract manufacturers association’s GRAS list. Among those of interest today is monomenthyl succinate (MMS) (FEMA# 3810) (18), which was later patented by Mane as a cooling agent for general use (20). In addition, menthol ethylene glycol carbonate (Frescolat[®] MGC), with FEMA# 3805, and menthol propylene glycol carbonate (Frescolat MPC), with FEMA# 3806, were first patented as tobacco flavorants (19), again to be later patented by Haarmann and Reimer for general cooling usages (21).

Q4
Q5

A number of other menthol-related cooling agents are commercially available: menthone glycerol ketal (Frescolat MGA) (22)—both the racemic (FEMA# 3808) and *leavo* forms (FEMA# 3807); the *leavo* form appears to be the main item of commerce. This material provides a clean cooling refreshing effect and as a partial replacement of peppermint oil has been shown to provide longer-lasting sweetness and a higher cooling sensation in chewing gum (23). (–)-Menthyl lactate (Frescolat ML) is faintly minty in odor and virtually tasteless with a pleasant, long-lasting cooling effect (24). Recently, Erman has shown that the (–)-ML of commerce has the ‘S’ configuration for the hydroxy moiety, indicating the fact that it is produced by the esterification of (–)-menthol with (S)-(+)-lactic acid (25). 3-(1-Menthoxy)propane-1,2-diol, known as MPD, Coolact[®] agent 10, TK-10, and coolant agent 10, is another important commercial cooling agent, which, in contrast to menthol, is essentially odorless (26). The cooling threshold (in mouth) is 1 ppm (about 20–100% that of menthol), and the time of cold-feeling maintenance is 20 to 25 minutes for a 100-ppm solution (about twice that of menthol). While the cooling strength of Coolact agent 10 is accepted as being about 20% to 25% that of menthol, it is also noted that “in a Vaseline ointment, 3-(1-menthoxy)propane-1,2-diol shows a cool feeling 2.0 to 2.5 times stronger than that of (–)-menthol” (27). The cool-feeling intensity of the (2S) isomer is 2 to 3 times that of the (2R) isomer and 1.5 to 2 times

233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290

superior to that of the racemic modification (28). Similarly, the related menthoxyalkanols, 3-(1-menthoxy)-2-methylpropane-1,2-diol (FEMA# 3849), 3-(1-menthoxy)ethanol (Coolact 5), FEMA# 4154, 3-(1-menthoxy)propan-1-ol, and 3-(1-menthoxy)butan-1-ol have cooling properties (29). Interestingly, cooling compounds such as 3-(1-menthoxy)propane-1,2-diol and 3-(1-menthoxy)-2-methylpropane-1,2-diol when admixed with warming sensates (e.g., vanillyl butyl ether, ginger extract, or capsicum tincture) provide increased warmth and longer-lasting warmth in cosmetic and flavor systems (27,30,31). Conversely, it has also been observed that admixtures of such cooling compounds with the warming sensate vanillin-MPD (the acetal of 3-(1-menthoxy)propane-1,2-diol and vanillin), FEMA# 3904, can increase the duration of cooling sensations (32) (Fig. 3).

(-)-Isopulegol (Coolact P), FEMA# 2962, having a chemical purity of better than 99.7% and an optical purity of not less than 99.7% ee, is odorless and gives a feeling of freshness, crispness, and coolness. The cooling strength is about 20% to 30% that of (-)-menthol (33). The *p*-menthane-3,8-diols (Coolact 38D, PMD38), FEMA# 4053, consist of a mixture of (+)-*cis* and




Figure 3 Menthoxy-related coolants.

291 (-)-*trans* PMD38 in a ratio of ~62:38 and possesses a cooling strength of about 11% that of
292 (-)-menthol (27,34). PMD38 is a nature identical material that occurs in a number of citronellal-
293 rich oils (e.g., *Litsea cubeba*, *Eucalyptus citriodora*) and is also effective as an insect repellent
294 (34–36). (-)-Monomenthyl glutarate (Physcool 2, MMG), FEMA# 4006, is a nature identical
295 cooling agent that has been found in *Litchi sinensis* accompanied by (-)-dimenthyl glutarate
296 (37). It has been described as “probably the longest-lasting oral cooling agent that is
297 commercially available” (38). Recently, an improved synthesis has been reported for both
298 MMG and MMS that minimizes the amount of diester impurities (39). Similarly, (-)-MMS has
299 been confirmed to be nature identical by its isolation from *Lycium barbarum* and *M. piperita* (37).
300 A recent description of MMS indicates that it is virtually tasteless and has well-balanced
301 cooling onset and length of cooling (38). Questice[®] (menthyl pyrrolidin-2-one 5-carboxylate)
302 was first patented as a composition of matter that acts as a long-lasting cooling and fresh
303 ingredient in toothpaste. The cooling properties are due to the enzymatic hydrolytic release of
304 menthol. A liquid form was produced by reacting (-)-menthol with L-pyrrolidin-2-one
305 carboxylic acid, while a crystalline form was produced when racemic DL-pyrrolidin-2-one
306 carboxylic acid is employed (40). Surprisingly, it did not appear on the GRAS list until 2005
307 with FEMA# 2155 (41). However, it has long been employed in various cosmetics, lotions, etc.
308 Recently, Erman has shown that the liquid form of Questice is a diastereoisomeric mixture of
309 (-)-menthyl 5-oxopyrrolidine-2-carboxylates with a ratio of the 5S:5R configuration of ~91:8,
310 while the solid form has a ratio of ~46:53 (25). (-)-Menthyl 3-hydroxybutyrate (MHB), FEMA#
311 4308, is another recent addition to the GRAS list (42). This is reported by workers at Takasago
312 as having a long-acting excellent cooling effect and is odorless and tasteless. Potential uses
313 include foods, drinks, cosmetics, pharmaceuticals, and cigarettes (43). Other workers indicate
314 that the cooling effect is slightly stronger than ML (about 48% the cooling strength of
315 menthol) (44). Firmenich workers have recently found that a diastereoisomeric mixture of the
316 6-isopropyl-3,9-dimethyl-1,4-dioxaspiro[4.5]decan-2-ones, prepared by reacting lactic acid
317 with *cis* and *trans*-menthones, provides a minty, fresh, *piperita*-type flavor that is remarkable
318 by its strength and cleanness. In combination with other cooling agents (e.g., menthyl succinate
319 or menthol), a synergist increase in cooling strength was found. In particular, the (3S,5R,6S,9R)
320 and (3S,5S,6S,9R) isomers are preferred (45). A patent describes the use of certain esters such as
321 (-)-menthyl methoxyacetate and (-)-menthyl 3,6-dioxaheptanoate as cooling agents (46). In
322 addition to the cooling properties, (-)-menthyl methoxyacetate has a head note and fruity taste
323 resembling that of menthyl acetate, whereas (-)-menthyl 3,6-dioxaheptanoate has a bitter taste.
324 Cubebol, a natural isolate of cubeb oil, in which it normally occurs at levels of 10% to 30% (47),
325 is a sesquiterpenoid alcohol that has a certain stereochemical resemblance to menthol and,
326 while not menthol derived, is included here for completeness. Cubebol has only a very weak
327 smell and taste and provides a refreshing effect that develops in the mouth after a delay of
328 approximately 1 to 2 minutes and lasts for approximately 30 minutes. It has applications in
329 flavors, oral care, pharmaceutical products, etc. (48).

330 Carboxamide Cooling Agents

331 During the early 1970s, Wilkinson Sword Ltd. conducted an extensive research program in
332 which they designed and evaluated about 1200 compounds for their cooling activity (49,50).
333 The interest in such compounds related to cooling agents without the minty and volatile
334 side effects of menthol, such as eye irritation, in aftershave lotions, etc. Over 25 U.S. patents
335 were issued on these materials (51). Of these original Wilkinson Sword compounds, three
336 were initially commercialized: WS-3 (*N*-ethyl-*p*-menthane-3-carboxamide) (52), WS-23
337 (2-isopropyl-*N*,2,3-trimethylbutyramide) (53), and WS-14 [*N*-([ethoxycarbonyl]methyl)-*p*-
338 menthane-3-carboxamide] (52). WS-3 was given GRAS status (FEMA# 3455) in 1975 (54) and
339 WS-23 (FEMA# 3804) in 1996 (55). Interestingly, WS-14 was used as a cooling agent for the
340 Northwind cigarette introduced into test market in 1981. This test market was short lived, but it
341 is not clear if this was because of market failure or concern that the additive testing conducted
342 was insufficient to pass Food and Drug Administration (FDA) scrutiny (56). WS-14 is
343 commercially available as ICE 4000 cooling sensate (57) and finds some applications as a
344 topical cooling agent (Fig. 4).

345 In 2007, WS-5 [ethyl 3-(*p*-menthane-3-carboxamido)acetate], which is currently the
346 coldest of all commercial cooling agents, was granted GRAS status as FEMA# 4309 (42). It has
347 been found that only highly purified WS-5 is suitable for flavoring purposes (58), as less pure
348

349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406




Figure 4 Successful Wilkinson sword coolants.

material exhibits a powerful bitter taste. WS-3 and WS-23 are currently the two largest volume carboxamide coolants. They are widely used in flavors, especially for chewing gum, breath fresheners, confectionaries, and oral care. They also find use in cosmetics (e.g., aftershave lotions). As both WS-3 and WS-23 are solids, there has been considerable interest in developing blends of such cooling agents that provide strong cooling but are easy to handle liquids. For example, it has been found that mixtures of ML, WS-3, and propylene glycol form stable liquid systems (59). It has also been shown that WS-3, WS-5, WS-14, and WS-23, alone or in certain combinations, when mixed with ML (or other coolants such as menthoxypropane-1,2-diol) will form stable liquid systems (60), and such mixtures often give a synergistic increase in cooling sensation. Similarly, eutectic mixtures of WS-3 and WS-23 provide liquid cooling systems (61,62), which can be used either as cooling agents or as flavor and saltiness enhancers.

Another compound that can be classified either as a carboxamide or a menthyl ester is *N,N*-dimethyl menthyl succinamide (FEMA# 4230 for the racemate). An IFF patent (63) describes this as having a cooling onset time of 25 seconds with cooling duration of 11.25 minutes. The taste/sensory profile is "cooling and refreshing on tongue, palate and front gums; fruity flavor with estery top-notes and sour undertones" (at 25 ppm in water). In a chewing gum at 0.2%, it increased sweetness and exhibited a pleasant and substantive cooling effect on the tongue and roof of the mouth.

Q6

Other examples of newly discovered carboxamides coolants are a series of analogs of WS-23 [such as *N*-(2-ethoxyethyl)-2-isopropyl-2,3-dimethylbutanamide] patented by Qaroma (64) and aryl carboxamide analogs (with the reversed amide configuration) by Givaudan (65), many with cooling intensities equal to or greater than WS-23. For example, *N*-(1-isopropyl-1,2-dimethylpropyl)-1,3-benzodioxole-5-carboxamide has about 2.2 times more cooling intensity as compared with 2 ppm of menthol (Fig. 5).




Figure 5 Recent WS-23 analogs.




Figure 6 New *p*-menthane carboxamide coolants.

Of particular interest are various aryl *p*-menthane-3-carboxamides, such as *N*-benzo[1,3]dioxol-5-yl-3-*p*-menthane-3-carboxamide and *N*-benzooxazol-4-yl-3-*p*-menthane-3-carboxamide, which are reported to have 100 times more cooling intensity than menthol (when compared with menthol at 2 ppm) (66).

In 2004, T. Hasegawa Co. Ltd. patented a new series of strong cooling compounds on the basis of alkyloxy amides of the *p*-menthane series, which exhibit no bitterness; compound HASE-1 is an example (Fig. 6) (67).

Further, Wei (68) has shown that several materials related to WS-5 possess strong cooling with remarkable cooling longevity. For example, the methyl and ethyl ester analogs of WS-5 (referred to as D-Ala-O-Me and D-Ala-O-Et, respectively) are produced from D-alanine (rather than glycine). Similarly, when D-homoserine lactone is employed, the resultant compound is *N*-(*R*)-2-oxotetrahydrofuran-3-yl-1-(1*R*,2*S*,5*R*)-*p*-menthane-3-carboxamide (referred to as "D-HSL"), which also is a potent long-lasting coolant. By combining suitable sympathomimetic amine drugs that act as α -adrenergic receptor agonists to form the corresponding *p*-menthane carboxamides, Wei found certain compounds (such as L-phenylephrine *p*-menthane carboxamide, referred to as CPS-195) that were effective as long-lasting coolants and possessed additional therapeutic properties (69). The cooling duration of a number of these, applied to the skin as a 1% wt/vol in a petrolatum-based ointment, versus leading coolants is shown in Figure 7.

In the last 10 years, there has been extensive patent activity relative to physiological cooling agents. Between 1998 and 2007, more than 280 patents were issued (25,63), and from January 2005 to December 2007 more than 300 patent applications have been filed. It is beyond the scope of this article to review all of these. However, it should be noted that a recent activity trend has been the patenting of various combinations of cooling agents, both to achieve improved cooling properties and/or for liquefaction of solid coolants (59,61,70,71). For example, it has been found that blends of menthyl glutarate with low levels of

457
458
459
460
461
462
463
464




Figure 7 Topical cooling duration 1% in ointment. Source: From Refs. 68, 69.

(-)-isopulegol and/or PMD38 exhibit a remarkable synergistic increase in cooling in oral-care products (71).

The relative "accepted" cooling strengths of important coolants are shown in Figure 8 (72,73).

It should be noted that "accepted" cooling strengths, primarily associated with topical skin cooling, do not always agree when compared to oral sensory panel results. This is clearly shown by results obtained by Wm. Wrigley Jr. Company sensory panels comparing 5% sucrose solutions of various coolants versus 100-ppm (-)-menthol, as shown in Figure 9 (71).




Figure 8 Approximate "accepted" cooling strengths versus menthol (as 100). Source: From Refs. 72, 73.




Figure 9 Relative oral cooling in 5% sucrose solutions versus 100 ppm menthol. Source: From Ref. 71.

COOLING COMPOUNDS AS INSECT REPELLENTS

As previously mentioned, the PMD38 have shown effectiveness as an insect repellent. Barnard has compared its efficacy against the leading insect repellants DEET, IR3535 [ethyl 3-(*N*-butyl-*N*-acetyl)-aminopropionate], and KBR3023 [sec-butyl 2-(2-hydroxyethyl)piperidine-1-carboxylate] (74,75).

Questice (menthyl pyrrolidone carboxylate) has also been patented as an insect repellent (76) and, Kalbe and Nentwig describe the use of ML or menthol glycerol acetal for repelling mites and other insects (77). Notably, Gautschi and Blondeau of Givaudan have discovered that WS-3 (*N*-ethyl-*p*-menthane-3-carboxamide) and related *N*-substituted *p*-menthane carboxamides have insect repelling activity against cockroaches equal to or exceeding that of DEET (diethyl-*m*-toluamide) (79). Another Givaudan patent application describes the use of a series of (-)-menthyl carbamates as insect repellents, but is silent relative to their cooling activity (80).

COLD RECEPTORS AND MECHANISM OF ACTION

The underlying process in thermoreception, whether hot or cold, is dependent on ion transport across cellular membranes. Cellular membranes consist of an oily phospholipid bilayer, which would be impermeable to ions such as K⁺ or Ca²⁺, except for receptor protein ion channels.

The flow of ions through these gated ion channels can cause rapid changes in ion concentrations, which in turn produce electrical signals that are the basis for many biological processes (80). In the case of thermoreceptors, these are activated when a thermal (or chemical) stimulus excites primary afferent sensory neurons of the dorsal or trigeminal ganglia (81).

In the last 12 years, there has been tremendous progress in determining the various receptor structural sequences. Thermoreceptors belong to the class of transient receptor potential (TRP) channels of which seven subfamilies exist (TRPC, TRPV, TRPM, TRPA, TRPP, TRPML, and TRPN). Six members of three TRP subfamilies are involved in mammalian

581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638

Table 4 Thermoreceptor Agonists

Chemical agonist (botanical source)	ThermoTRP
Capsaicin (hot chilli peppers, e.g., Tabasco®)	TRPV1
Piperine (black pepper corns)	TRPV1
Allicin (fresh garlic)	TRPV1, TRPA1
Camphor (<i>Cinnamomum camphora</i>)	TRPV3, TRPV1
Δ -9-Tetrahydrocannabinol (<i>Cannabis sativa</i>)	TRPV2, TRPA1
2-Aminoethoxydiphenyl borate (synthetic)	TRPV1, TRPV2, TRPV3
4- α -phorbol 12,13-didecanoate (synthetic)	TRPV4
(-)-Menthol (peppermint)	TRMP8, TRPV3
1,8-Cineole, eucalyptol (eucalyptus)	TRPM8
WS-3 (synthetic)	TRPM8
Icilin (synthetic)	TRPM8, TRPA1
Cinnamaldehyde (cinnamon, cassia)	TRPA1, TRPV3
Allyl isothiocyanate (mustard, horseradish)	TRPA1
Benzyl isothiocyanate (mustard, horseradish)	TRPA1
Phenethyl isothiocyanate (mustard, horseradish)	TRPA1

Abbreviation: TRP, transient receptor potential.
Source: From Refs. 80, 84.

temperature-sensitive thermoreception. The closely related TRPV analogs are activated by heat, TRPV1 ($\geq 43^\circ\text{C}$), TRPV2 ($\geq 52^\circ\text{C}$), TRPV3 (22–40°C), and TRPV4 ($> \sim 27^\circ\text{C}$), while TRPM8 ($< \sim 28^\circ\text{C}$) and TRPA1 ($< \sim 18^\circ\text{C}$) are activated by cold (80). Certain types of chemical agonists activate these same thermoTRP channels. TRPV1 was the first thermoreceptor characterized and is referred to as a vanilloid receptor, as it is activated by capsaicin as well as heat. The cold and menthol receptor, TRPM8, was characterized by McKemy, Neuhausser, and Julius (82) and by Peier et al. (83) in 2002. Paradoxically, the cold receptor TRPA1, which is activated by noxious cold to produce a pain-like sensation, produces a human sensorial effect often described as “hot.”

Table 4 provides examples of chemical agonists that activate these thermoTRPs.

All of these thermoTRPs are gated Ca^{2+} channels consisting of six transmembrane domains (TM1–TM6) flanked by large N- and C-terminal cytoplasmic domains (80). A schematic representation is shown in Figure 10 with the putative ion channel between TM5–TM6 in TRPM8, which is activated by menthol and other cold stimuli. In the case of TRP channels, it has been shown that they can oligomerize into tetramer assemblies, which presumably modulate the calcium ion gating processes (85–87).

Much of the knowledge gained on TRP activation by chemical stimuli has been derived by genetic expression of putative receptor domains and measurement of Ca^{2+} flux intensity by fluorometric imaging assays. Behrendt et al. used this technique to screen 70 odorants and menthol-related substances for activity on the recombinant cold-menthol receptor TRPM8




Figure 10 TRPM8 receptor channel.




Figure 11 Efficacy of coolants on the TRPM8 receptor (by Ca²⁺ fluorometric assay). *Source:* From Refs. 72, 73, 88.

(mTRPM8), as expressed in HEK293 cells (88). The percentage efficacy of the most active candidates as compared to menthol is shown in Figure 11.

Although the fluorometric assay technique does not always translate into the human perception scale, it is already being used in industry to screen for promising new coolants (89).

In conclusion, from peppermint to menthol and to a plethora of new novel cooling compounds, we are now beginning to understand the importance of cooling substances even in the genetics of life. From early menthol-camphor-based over-the-counter (OTC) pharmaceuticals, which created famous trademarks such as Vicks[®] Vaporub and Mentholatum[®] in the early 20th century, to improved modern toothpastes, gums, breath fresheners, and cosmetic lotions, we expect this is just the beginning of even greater things for the future.

REFERENCES^a

Q7

1. Guenther E. The Essential Oils, Vol. 3. Huntington NY: Robert E Krieger Pub. Co., 1974:640–676.
2. Gildemeister E, Hoffmann Fr. The Volatile Oils. English translation by E. Kremers. Milwaukee: Pharmaceutical Review Pub. Co., 1900:631–640.
3. Flückiger FA. Pharmakognosie des Pflanzenreiches, 2d ed. Berlin: R. Gaertner’s Buchhandlung, 1883: 686. Availabel at: <http://www.digibib.tu-bs.de/?docid=00000684>. Accessed December 4, 2007.
4. Clark GS. Aroma chemical profile – menthol. Perfumer & Flavorist 2007; 32(12):32–47.
5. Menthol—A Cool Place. Availabel at: <http://www.leffingwell.com/menthol1/menthol1.htm>. Accessed December 5, 2007.
6. Hopp R, Lawrence BM. Natural and synthetic menthol. In: Lawrence BM, ed. Mint: The Genus Mentha, Boca Raton, FL: CRC Press, 2006: 371–398.
7. Fleischer J, Bauer K, Hopp R. Separating Optically Pure d- and l-Isomers of Menthol, Neomenthol and Isomenthol. US patent 3 943 181. March 9, 1976.
8. Menthol—A Cool Place. Availabel at: <http://www.leffingwell.com/menthol10/menthol10.htm>. Accessed December 5, 2007.

Q8

^a Note on Trademarks: Words that we know or have reason to believe constitute registered trademarks, [®], are designated as such in the text. However, neither the presence nor absence of such designation should be regarded as affecting the legal status of any trademark.

- 697 9. Akutagawa S. Enantioselective isomerization of allylamine to enamine: practical asymmetric
698 synthesis of (–)-menthol by Rh-BINAP catalysts. *Top Catal* 1997; 4(3–4):271–274.
- 699 10. Tani T, Yamagata T, Akutagawa S, et al. Metal-assisted terpenoid synthesis. 7. Highly enantioselective
700 isomerization of prochiral allylamines catalyzed by chiral diphosphine rhodium (I) complexes.
701 Preparation of optically active enamines. *J Am Chem Soc* 1984; 106(18):5208–5217.
- 702 11. Noyori R. Asymmetric catalysis: science and opportunities (Nobel Lecture, December 8, 2001) in *Les*
703 *Prix Nobel. The Nobel Prizes 2001*. Frängsmyr T. ed. Stockholm: Nobel Foundation, 2002: 192–193.
704 Available at: http://nobelprize.org/nobel_prizes/chemistry/laureates/2001/noyori-lecture.pdf. Accessed
705 December 5, 2007.
- 706 12. Read J. Recent progress in the menthone chemistry. *Chem Rev* 1930; 7(1):1–50.
- 707 13. Read J, Grubb WJ, Malcolm D. Researches in the menthone series. Part XI. Diagnosis and characterization
708 of the stereoisomeric menthols. *J Chem Soc* 1933; 170–173.
- 709 14. Hopp R. Menthol: its origins, chemistry, physiology and toxicological properties. *Rec Adv Tob Sci*
710 1993; 19:3–46.
- 711 15. Leffingwell JC, Shackelford RE. Leavo menthol – syntheses and organoleptic properties. *Cosmetics*
712 *and Perfumery* 1974; 89(6):69–89.
- 713 16. Ayya N, de Wijk R, Frijters J. Sensory characteristics of various menthols using an injection olfactometer.
714 1994. Available at: <http://legacy.library.ucsf.edu/tid/wzg93f00>. Accessed December 4, 2007.
- 715 17. Milleville G, Winkler G. Verfahren zum Mentholisieren von Tabak und zur Herstellung von
716 Mentholzigaretten. German patent 1 065 767. September 18, 1959.
- 717 18. Jarboe CH. Smoking tobacco product and method of making the same. US patent 3 111 127.
718 November 19, 1963.
- 719 19. Mold JD, Kallianos AG, Shellburne FA. Tobacco incorporating carbonate esters of flavorants. US
720 patent 3 332 428. July 25, 1967.
- 721 20. Mane JM. Coolant compositions. US patent 5 725 865. March 19, 1998.
- 722 21. Pelzer R, Surburg H, Hopp R. Mittel mit physiologischem Kühleffekt und für diese Mittel geeignete
723 wirksame Verbindungen. German patent 4 226 043. November 2, 1994.
- 724 22. Grub H, Pelzer R, Hopp R, et al. Compositions which have a physiological cooling effect, and active
725 compounds suitable for these compositions. US patent 5 266 592. November 30, 1993.
- 726 23. Greenberg MJ. Enhanced flavors using menthone ketals. US patent 5 348 750. September 20, 1994.
- 727 24. Bauer K, Bruening J, Grueb H. Mittel mit physiologischer Kuhlwirkung. German patent 2 608 226.
728 September 8, 1977.
- 729 25. Erman M. New developments in physiological cooling agents. *Perfumer & Flavorist* 2007; 32(10):20–35.
- 730 26. Amaro A, Moroe M, Yoshida T. 3-Levo-Menthoxyp propane-1,2-diol. US patent 4 459 425. July 10, 1984.
- 731 27. Shiroyama K, Sawano K, Ohta H. Cool feeling composition. US patent 6 328 982. Dec 11, 2001.
- 732 28. Amano A, Tokoro K. (2S)-3-(1R, 2S, 5R)-5-methyl-2-(1-methylethyl)-cyclohexyl[oxy]-1,2-propanediol,
733 process for producing the same, and compositions containing the same. US patent 5 608 119. March 4, 1997.
- 734 29. Green C, Nakatsu T, Ishizaki T. (1R, 2S, 5R)-3-1-menthoxyalkan-1-ol cooling sensate. US patent 6 956
735 139. October 18, 2005.
- 736 30. Kumamoto H, Ohta H. Warming composition. US patent 6 673 844. January 6, 2004.
- 737 31. Kumamoto H, Kitamura T. Warming composition for food and drink or for oral care preparation. US
738 patent 6 838 106. January 4, 2005.
- 739 32. Nakatsu T, Green CB, Reitz GA. Fragrance composition containing 4-(1-menthoxymethyl)-2-phenyl-
740 1,3-dioxolane or its derivatives. US patent 5 753 609. May 19, 1998.
- 741 33. Yamamoto T. Method for purifying (–)-N-isopulegol and citrus perfume composition containing
742 (–)-N-isopulegol obtained by the method. US patent 5 773 410. June 30, 1998.
- 743 34. Kenmochi H, Akiyama T, Yuasa Y, et al. Method for producing para-menthane-3,8-diol. US patent 5
744 959 161. September 28, 1999.
- 745 35. Beldock DT, Beldock JA, Mudge G. Insect repellent blends, lotions, and sprays. US patent 5 621 013.
746 April 15, 1997.
- 747 36. West Nile Virus. Available at: <http://www.cdc.gov/ncidod/dvbid/westnile/RepellentUpdates.htm>.
748 Accessed December 16, 2007.
- 749 37. Hiserodt RD, Adedeji J, John TV. et al. Identification of monomenthyl succinate, monomenthyl
750 glutarate, and dimethyl glutarate in nature by high performance liquid chromatography-tandem
751 mass spectrometry. *J Agric Food Chem* 2004; 52(11):3536–3541.
- 752 38. Dewis ML. Molecules of taste and sensation. In: Rowe DJ, ed. *Chemistry and Technology of Flavors*
753 *and Fragrances*, Oxford: Blackwell Pub. Ltd, 2004:199–243.
- 754 39. Erman MB, Snow JW, Lebedev MY. Process for making monomenthyl esters. US patent 7 247 743. July
24, 2007.
40. Humbert FEL, Guth G. Esters of menthol and 2-pyrrolidone or piperidine carboxylic acids. US patent
3 917 613. November 4, 1975.
41. Smith RL, Cohen SM, Doull J, et al. GRAS Flavoring substances 22. *Food Technol* 2005; 59(8): 24–28,
31–32, 34, 36–62.

755 42. Waddell WJ, Cohen SM, Feron VJ, et al. GRAS flavoring substances 23. *Food Technol* 2007; 61(8):
756 22-48.
757 43. Yamamoto T, Amano A, Kobayashi T. L-Menthyl 3-hydroxybutyrate, production thereof, and chilling
758 agent containing said compound as active component. Japanese patent 6 119 4049. August 28, 1986.
759 44. Pringle S, Brassington D. The chemistry of F&F: physiological coolants. *Perfumer & Flavorist* 2007;
760 32(1):38-42.
761 45. Giersch WK, Vanrietvelde CLA, et al. Flavor ingredients for cooling preparations, US patent
762 application 20060249167. November 9, 2006.
763 46. Frerot E, Van Beem N. Compounds derived from menthol and use as refreshing agent. US patent 6
764 359 168. March 19, 2002.
765 47. Boelens Aroma Chemical Information Service, ESO 2000 (update 2006) – The Complete Database of
766 Essential Oils. Canton GA: Leffingwell & Associates, Pub. 2006.
767 48. Velazco MI, Wuensche L, Deladoey P. Use of cubebol as a flavoring ingredient. US patent 6 214 788.
768 April 10, 2001.
769 49. Watson HR. Flavor characteristics of synthetic cooling compounds. In: Apt CM, ed. *Flavor: It's*
770 *Chemical, Behavioral & Commercial Aspects*. Boulder: Westview Press, 1977:31-50.
771 50. Watson HR, Hems R, Roswell DG, et al. New compounds with the menthol cooling effect. *J Soc*
772 *Cosmet Chem* 1978; 29(4): 185-200.
773 51. The US Patent and Trademark Office. Available at: <http://www.uspto.gov/patft/index.html>.
774 52. Watson HR, Roswell DG, Spring DJ. N-Substituted paramenthane carboxamides. US patent 4 178 459.
775 December 11, 1979.
776 53. Roswell DG, Spring D, Hems R. Aliphatic N-Substituted Tertiary Amides Possessing Physiological
777 Cooling Activity. Great Britain patent 1 421 744. January 21, 1976.
778 54. Oser BL, Ford RA. Recent progress in the consideration of flavoring ingredients under the Food
779 Additives Amendment. 9. GRAS substances. *Food Technol* 1975; 29(8):70-72.
780 55. Smith RL, Newberne P, Adams TB, et al. GRAS flavoring substances 17. *Food Technol* 1996; 50(10):72-81.
781 56. Legacy Tobacco Documents. Available at: <http://legacy.library.ucsf.edu/tid/mds88d00>. Accessed
782 December 18, 2007.
783 57. Qaroma. Available at: <http://www.qaroma.com>. Accessed December 18, 2007.
784 58. Erman MB, Whelan P. Physiological cooling compositions containing highly purified ethyl ester of
785 N-5-methyl-2-(1-methylethyl) cyclohexyl. carbonyl]glycine. US patent 7 189 760. March 13, 2007.
786 59. Galopin C, Moraes E, Tigani L. Coolant solutions and compositions comprising the same. US patent
787 application 20060051301. March 9, 2006.
788 60. Erman MB, Whelan PJ, Snow JW. Physiological cooling compositions. US patent application
789 20050265930. December 1, 2005.
790 61. Moza AK, Sun H, Leffingwell JC. Liquid composition of 2-Isopropyl-N,2,3-trimethylbutyramide and
791 N-Ethyl-p-menthane-3-carboxamide, its preparation method and its applications as a cooling agent
792 and flavor enhancer. US patent application 20070048424. March 1, 2007.
793 62. Moza AK, Rohde SE, Leffingwell JC. Cooling agents as flavor and saltiness enhancers. US patent
794 application 20070059417. March 15, 2007.
795 63. Dewis ML, Huber ME, Cossette MV. Menthyl half acid ester derivatives, processes for preparing
796 same, and uses thereof for their cooling/refreshing effect in consumable materials. US patent 6 884
797 906. April 26, 2005.
798 64. Sun H. Novel compounds with physiological cooling effect. US patent 7 030 273. April 18, 2007.
799 65. Cole L, Furrer SM, Galopin C. Cooling Compounds. PCT WO 2006/099762. September 28, 2006.
800 66. Cole L, Furrer SM, Galopin C. Menthane Carboxamide Derivatives Having Cooling Properties. PCT
801 WO 2006/092074. September 8, 2006.
802 67. Takazawa O, Wanatabe H, Iso M. p-Menthane Derivative and Cold Sensing Agent Containing the
803 Same. Japanese patent 2004059474. February 26, 2004.
804 68. Wei ET. N-Alkylcarbonyl-Amino Acid Ester and N-Alkylcarbonyl-Amino Lactone Compounds and
805 Their Use. PCT WO 2006103401. October 5, 2006.
806 69. Wei ET. N-arylshydroxyalkylidene-carboxamide compositions and methods. US patent application
807 20070155755. July 5, 2007.
808 70. Kiefer J, Harvey JE. Cooling compositions. US patent application 20070077331. April 5, 2007.
809 71. Johnson SS, Stawski BZ, Sheldon GT. Confections Containing a Blend of Physiological Cooling
810 Agents. US patent application 20070248717. October 25, 2007.
811 72. Erman M. Progress in physiological cooling agents. *Perfumer & Flavorist* 2004; 29(8):34-50.
812 73. Leffingwell JC. Cool without Menthol & Cooler than Menthol and Cooling Compounds as Insect
Repellents. Available at: http://www.leffingwell.com/cooler_than_menthol.htm. Accessed December
18, 2007.
74. Barnard DR. Global collaboration for development of pesticides for public health (GCDPP):
Repellents and toxicants for personal protection. Geneva Switzerland:WHO Position Paper, World
Health Organization, 2000; 1-49.

- 813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
75. Barnard, DR, Bernier UR, Posey KH, et al. Repellency of IR3535, KBR3023, para-menthane-3,8-diol, and Deet to Black Salt Marsh Mosquitoes (*Diptera: Culicidae*) in the Everglades National Park USA J Med Entomol 2002; 39(6):895–899.
 76. Watkins S, Hills MJ, Birch RA. Use of menthyl-2 pyrrolidone-5-carboxylate as an insect repellent. US patent 6 451 844. September 17, 2002.
 77. Kalbe J, Nentwig G. Non-toxic, inexpensive arthropodicide comprises diacid ester, N-methyl-pyrrolidone, caprolactone or glycerol or menthol derivative, effective against insects, mites and ticks, especially dust mites. German patent 19840321. March 9, 2000.
 78. Blondeau P, Gautschi M. Composition having insect repellent characteristics. US patent application 20040028714. February 12, 2004.
 79. Gautschi M. Insect repellents. US patent application 20060063764. March 23, 2006.
 80. Voets T, Talavera K, Owsianik G, et al. Sensing with TRP channels. Nature Chem Biol 2004; 1(2):85–92.
 81. Jordt S-E, McKemy DD, Julius D. Lessons from peppers and peppermint: the molecular logic of thermosensation. Curr Opin Neurobiol 2003; 13(4):487–492.
 82. McKemy DD, Neuhäusser WM, Julius D. Identification of a cold receptor reveals a general role for TRP channels in thermosensation. Nature 2002; 416:52–58.
 83. Peier AM, Moqrich A, Hergarden AC, et al. A TRP channel that senses cold stimuli and menthol. Cell 2002; 108(5):705–715.
 84. Jordt S-E, Bautista DM, Chuang H, et al. Mustard oils and cannabinoids excite sensory nerve fibres through the TRP channel ANKTM1. Nature 2004; 427:260–265.
 85. Hoenderop JGJ, Voets T, Hoefs S, et al. Homo- and heterotetrameric architecture of the epithelial Ca²⁺ channels TRPV5 and TRPV6. EMBO J 2003; 22(4):776–785.
 86. Voets T, Janssens A, Droogmans G, et al. Outer pore architecture of a Ca²⁺-selective TRP channel. J Biol Chem 2004; 279(15):15223–15230.
 87. Phelps CB, Gaudet R. The role of the N Terminus and Transmembrane Domain of TRPM8 in Channel Localization and Tetramerization. J Biol Chem 2007; 282(50):36474–36480.
 88. Behrendt H-J, Germann T, Gillen C, et al. Characterization of the mouse cold-menthol receptor TRPM8 and vanilloid receptor type-1 VR1 using a fluorometric imaging plate reader (FLIPR) assay. Brit J Pharm 2004; 141(4):737–745.
 89. Servant GS, Brust P, Moyer B, et al. Optimized TRPM8 nucleic acid sequences and their USE in cell based assays and test kits to identify TRPM8 Modulators. US patent application 20070259354. November 8, 2007.

BOOK: H6963_Barel		CHAPTER 65
TO: CORRESPONDING AUTHOR AUTHOR QUERIES - TO BE ANSWERED BY THE AUTHOR		
The following queries have arisen during the typesetting of your manuscript. Please answer these queries.		
Q1	Page: 2 AU: Please provide the reference for "Haarmann and Reimer" and add to the Ref. list.	
Q2	Page: 2 AU: Please check whether this should be Ref. no. 4, viz., "Clark GS. Aroma chemical profile — menthol. Perfumer & Flavorist 2007; 32(12): 32-47."	
Q3	Page: 2 AU: Please check whether this should be Ref. no. 4, viz., "Clark GS. Aroma chemical profile — menthol. Perfumer & Flavorist 2007; 32(12): 32-47."	
Q4	Page: 4 AU: Please provide expansion for "GRAS."	
Q5	Page: 4 AU: Please provide expansion for "FEMA," if necessary.	
Q6	Page: 7 AU: Please provide expansion for "IFF," if necessary.	
Q7	Page: 12 AU: Please note that the note on trademarks has been converted into footnote as per style.	
Q8	Page: 12 AU: Please check for accuracy information inserted in Refs. 5, 8, 36, 51, 56, and 57.	
Q9	Page: 2 AU: OK to insert "Product category and Menthol %" as column heads 1 and 2, respectively, of Table 2?	
Q10	Page: 3 AU: FIGURE TO BE CORRECTED: THE FIRST MENTHOL SHOULD BE (+)-MENTHOL. AU: Please note that the preceding query is by M Paye.	
Q11	Page: 2 ED: We have kept Table 1 & Table 4 body text in light face for consistency throughout the chapter. Is it Ok? Please confirm.	

Chapter: 65

Contributors : John C. Leffingwell¹

Affiliation : ¹Leffingwell & Associates, Canton, Georgia, USA

Address : 4699 Arbor Hill Road, Canton, Georgia 30115

Phone : 1 770 889 5111

Fax : 1 770 887 0089

Email : leffingwell@mindspring.com

Lead Author : John C. Leffingwell